

GAPDH(HRP conjugated) Rabbit mAb

Catalog No: #49380

Package Size: #49380-1 50ul #49380-2 100ul



SAB[®]
Signalway Antibody

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com


Description

Product Name	GAPDH(HRP conjugated) Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	JF81-04
Purification	ProA affinity purified
Applications	WB
Species Reactivity	Hu, Ms, Rt, Monkey, Chicken, Fish
Immunogen Description	recombinant protein
Conjugates	HRP
Other Names	38 kDa BFA-dependent ADP-ribosylation substrate antibody aging associated gene 9 protein antibody Aging-associated gene 9 protein antibody BARS-38 antibody cb609 antibody EC 1.2.1.12 antibody Epididymis secretory sperm binding protein Li 162eP antibody G3P_HUMAN antibody G3PD antibody G3PDH antibody GAPD antibody GAPDH antibody Glyceraldehyde 3 phosphate dehydrogenase antibody Glyceraldehyde-3-phosphate dehydrogenase antibody HEL-S-162eP antibody KNC-NDS6 antibody MGC102544 antibody MGC102546 antibody MGC103190 antibody MGC103191 antibody MGC105239 antibody MGC127711 antibody MGC88685 antibody OCAS, p38 component antibody OCT1 coactivator in S phase, 38-KD component antibody peptidyl cysteine S nitrosylase GAPDH antibody Peptidyl-cysteine S-nitrosylase GAPDH antibody wu:fb33a10 antibody
Accession No.	Swiss-Prot#:P04406
Calculated MW	36 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C

Application Details

WB: 1:500-1:1,000

Images



Western blot analysis of GAPDH on different lysates using anti-GAPDH antibody at 1/1,000 dilution. Positive control:
Lane 1: Hela
Lane 2: PC-12
Lane 3: NIH/3T3

Background

Glyceraldehyde-3-phosphate dehydrogenase (GAPDH), also called uracil DNA glycosylase, catalyzes the reversible oxidative phosphorylation of glyceraldehyde-3-phosphate in the presence of inorganic phosphate and nicotinamide adenine dinucleotide (NAD), an important energy-yielding step in carbohydrate metabolism. While GAPDH has long been recognized as playing an integral role in glycolysis, additional functions of GAPDH include acting as a uracil DNA glycosylase, activating transcription, binding RNA and involvement in nuclear RNA export, DNA replication and DNA repair. Expression of GAPDH is upregulated in liver, lung and prostate cancers. GAPDH translocates to the nucleus during apoptosis. GAPDH complexes with neuronal proteins implicated in human neuro-degenerative disorders including the β -Amyloid precursor, Huntingtin and other triplet repeat neuronal disorder proteins.

References

Note: This product is for in vitro research use only and is not intended for use in humans or animals.