JAK2 Rabbit mAb

Catalog No: #58700

Package Size: #58700-1 50ul #58700-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


Description

| Product Name | JAK2 Rabbit mAb |
|-----------------------|--|
| Host Species | Rabbit |
| Clonality | Monoclonal |
| Isotype | Rabbit IgG |
| Purification | Affinity-chromatography |
| Applications | WB ICC/IF IP |
| Species Reactivity | Human Mouse Rat |
| Specificity | JAK2 Antibody detects endogenous levels of total JAK2 |
| Immunogen Description | A synthesized peptide derived from human JAK2 |
| Other Names | EC 2.7.10.2; JAK-2; JAK2; Janus kinase 2; kinase Jak2; JTK10; THCYT3; |
| Accession No. | Uniprot:O60674 |
| Formulation | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Storage | Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |


Application Details

WB 1:500~1:2000 ICC/IF 1:50~1:200 IP 1:50-100

Images



Western blot analysis of JAK2 expression in Jurkat cell lysate.



Immunofluorescent analysis of Jurkat cells, using JAK2 Antibody .

Background

Phosphorylated STATs then form homodimer or heterodimers and translocate to the nucleus to activate gene transcription. For example, cell stimulation with erythropoietin (EPO) during erythropoiesis leads to JAK2 autophosphorylation, activation, and its association with erythropoietin receptor (EPOR) that becomes phosphorylated in its cytoplasmic domain. Then, STAT5 (STAT5A or STAT5B) is recruited, phosphorylated and activated by JAK2.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.